

Neutrino Oscillations With Three Active and Three Sterile Neutrinos

Leonard S. Kisslinger

Department of Physics, Carnegie Mellon University, Pittsburgh PA 15213

September 14, 2015

PACS Indices:11.30.Er,14.60.Lm,13.15.+g

Keywords: sterile neutrinos, neutrino oscillations, U-matrix

Abstract

This is an extension of estimates of the probability of μ to e neutrino oscillation with one sterile neutrino to three sterile neutrinos, using a 6x6 matrix. Since the mixing angle for only one sterile neutrino has been experimentally determined, we estimate the μ to e neutrino oscillation probability with different mixing angles for two of the sterile neutrinos.

1 Introduction

Most recent theories of neutrino oscillations have used a 3x3 S-matrix approach with three active neutrinos[1, 2, 3]. Recent experiments on neutrino oscillations[4] have suggested the existence of at least one sterile neutrino with the mass and mixing angles used in the present work. See Ref[4] for references to earlier experiments, and Refs[5, 6] for reviews of sterile neutrino oscillations with references to experimental and theoretical publications.

In the present work we use a U-matrix approach, introduced for active neutrinos with a 3x3 U-matrix[7], and extended to a 4x4 U-matrix with one sterile neutrino in a recent study of $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$, the transition probability for a muon neutrino to oscillate to an electron neutrino[8, 9]. We introduce a 6x6 U-matrix for three active and three sterile neutrinos, an extension of previous work with six neutrinos[10]. An early study of the effect of adding 3 sterile neutrinos may be found in Ref[11], where it was found that in a broad class of theories consistent with grand unification, the neutrino mixing angles are likely to be comparable to the corresponding quark mixing angles and

might be much larger in a special case. This result holds for a wide range of mass ratios for the light-neutrino Majorana masses.

2 6x6 U-Matrix

Active neutrinos with flavors ν_e, ν_μ, ν_τ and three sterile neutrinos, $\nu_{s_1}, \nu_{s_2}, \nu_{s_3}$ are related to neutrinos with definite mass by

$$\nu_f = U \nu_m, \quad (1)$$

where U is a 6x6 matrix and ν_f, ν_m are 6x1 column vectors. We use the notation $s_{ij}, c_{ij} = \sin\theta_{ij}, \cos\theta_{ij}$, with $\theta_{12}, \theta_{23}, \theta_{13}$ the mixing angles for active neutrinos; and $s_\alpha = \sin(\alpha), c_\alpha = \cos(\alpha), s_\beta = \sin(\beta)$, etc, where α, β, γ are sterile-active neutrino mixing angles.

$$U = O^{23} O^{13} O^{12} O^{14} O^{24} O^{34} O^{15} O^{25} O^{35} O^{45} O^{16} O^{26} O^{36} O^{46} O^{56} \quad (2)$$

with (O^{45}, O^{46} , and O^{56} , giving sterile-sterile neutrino mixing, are not shown)

$$\begin{aligned} O^{23} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & c_{23} & s_{23} & 0 & 0 & 0 \\ 0 & -s_{23} & c_{23} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & O^{13} &= \begin{pmatrix} c_{13} & 0 & s_{13} & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ -s_{13} & 0 & c_{13} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \\ O^{12} &= \begin{pmatrix} c_{12} & s_{12} & 0 & 0 & 0 & 0 \\ -s_{12} & c_{12} & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & O^{14} &= \begin{pmatrix} c_\alpha & 0 & 0 & s_\alpha & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ -s_\alpha & 0 & 0 & c_\alpha & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \\ O^{24} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & c_\alpha & 0 & s_\alpha & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & -s_\alpha & 0 & c_\alpha & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & O^{34} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & c_\alpha & s_\alpha & 0 & 0 \\ 0 & 0 & -s_\alpha & c_\alpha & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
O^{15} &= \begin{pmatrix} c_\beta & 0 & 0 & 0 & s_\beta & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ -s_\beta & 0 & 0 & 0 & c_\beta & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & O^{25} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & c_\beta & 0 & 0 & s_\beta & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & -s_\beta & 0 & 0 & c_\beta & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \\
O^{35} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & c_\beta & 0 & s_\beta & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -s_\beta & 0 & c_\beta & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & O^{16} &= \begin{pmatrix} c_\gamma & 0 & 0 & 0 & 0 & s_\gamma \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ -s_\gamma & 0 & 0 & 0 & 0 & c_\gamma \end{pmatrix} \\
O^{26} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & c_\gamma & 0 & 0 & 0 & s_\gamma \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & -s_\gamma & 0 & 0 & 0 & c_\gamma \end{pmatrix} & O^{36} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & c_\gamma & 0 & 0 & s_\gamma \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -s_\gamma & 0 & 0 & c_\gamma \end{pmatrix}
\end{aligned}$$

$\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ is obtained from the 6x6 U matrix and the neutrino mass differences $\delta m_{ij}^2 = m_i^2 - m_j^2$ for a neutrino beam with energy E and baseline L by

$$\mathcal{P}(\nu_\mu \rightarrow \nu_e) = \text{Re} \left[\sum_{i=1}^6 \sum_{j=1}^6 U_{1i} U_{1j}^* U_{2i}^* U_{2j} e^{-i(\delta m_{ij}^2/E)L} \right], \quad (3)$$

an extension the 4x4[8, 9] theory with one sterile neutrino, which used the 3x3 formalism of Ref[7], to a 6x6 matrix formalism[10]. From Eq(2), multiplying the 12 6x6 O matrices, we obtain the matrix U. With $\delta_{CP}=0$, $U_{ij}^* = U_{ij}$, so we only need U_{1j}, U_{2j} .

$$\begin{aligned}
U_{11} &= .821ca \, cb \, cg \\
U_{12} &= cg((.554ca - .821sa^2)cb - .821ca \, sb^2) - .821ca \, cb \, sg^2 \\
U_{13} &= cg((.15ca - .554sa^2 - .821ca \, sa^2)cb - (.554ca - .821sa^2)sb^2 \\
&\quad - .821ca \, cb \, sb^2) - .821ca \, cb \, cg \, sg^2 - ((.554ca - .821sa^2)cb - .821ca \, sb^2)sg^2 \\
U_{14} &= cg(cb(.15sa + .554ca \, sa + .821ca^2 \, sa) - .821ca \, cb^2 \, sb^2 \\
&\quad - (.554ca - .821sa^2)cb \, sb^2 - (.15ca - .554sa^2 - .821cas a^2)sb^2) - .821ca \, cb \, sg^2 cg^2 \\
&\quad - cg((.554ca - .821sa^2)cb - .821ca \, sb^2)sg^2 - (cb(.15ca - .554sa^2 - .821cas a^2) \\
&\quad - .821ca \, cb \, sb^2 - (.554ca - .821sa^2)sb^2)sg^2
\end{aligned}$$

$$\begin{aligned}
U_{15} &= cg(.821ca\ sb\ cb^3 + (.15sa + .554ca\ sa + .821ca^2\ sa)sb \\
&\quad + (.554ca - .821sa^2)cb^2\ sb + (.15ca - .554sa^2 - .821ca\ sa^2)cb\ sb) \\
&\quad - .821ca\ cb\ cg^3sg^2 - cg^2(cb(.554ca - .821sa^2) - .821sb^2)sg^2 \\
&\quad - cg(cb(.15ca - .554sa^2 - .821ca\ sa^2) - .821ca\ cb\ sb^2 \\
&\quad - (.554ca - .821sa^2)sb^2sg^2 - (cb(.15sa + .554ca\ sa + .821ca^2\ sa) - .821ca\ cb^2\ sb^2 \\
&\quad - cb(.554ca - .821sa^2)sb^2 + (.15ca - .554sa^2 - .821ca\ sa^2)sb^2)sg^2 \\
U_{16} &= .821ca\ cb\ sg\ cg^4 + (.821ca\ cb^3\ sb + (.15sa + .554ca\ sa + .821ca^2\ sa)sb \\
&\quad + cb^2(.554ca - .821sa^2)sb + cb(.15ca - .554sa^2 - .821ca\ sa^2)sb)sg \\
&\quad + cg^3((.554ca - .821sa^2)cb - .821ca\ sb^2)sg + \\
&\quad cg^2(cb(.15ca - .554sa^2 - .821ca\ sa^2) - .821ca\ cb\ sb^2 \\
&\quad - (.554ca - .821sa^2)sb^2)sg \\
&\quad + cg(cb(.15sa + .554ca\ sa + .821ca^2\ sa) - .821ca\ cb^2\ sb^2 \\
&\quad - cb(.554ca - .821sa^2)sb^2 - (.15ca - .554sa^2 - .821ca\ sa^2)sb^2)sg, \tag{4}
\end{aligned}$$

$$\begin{aligned}
U_{21} &= -.484ca\ cb\ cg \\
U_{22} &= cg(.527ca + .484sa^2)cb - .821ca\ sb^2) + .484ca\ cb\ sg^2 \\
U_{23} &= cg((.699ca - .527sa^2 + .484ca\ sa^2)cb - (.527ca + .484sa^2)sb^2 + .484ca\ cb\ sb^2) \\
&\quad + .484ca\ cb\ cg\ sg^2 - ((.527ca + .484sa^2)cb + .484ca\ sb^2) * sg^2 \\
U_{24} &= cg(cb(.699sa + .527ca\ sa - .484ca^2\ sa) + .484ca\ cb^2\ sb^2 \\
&\quad - (.527ca + .484sa^2)cb\ sb^2 - (.699ca - .527sa^2 + .484ca\ sa^2)sb^2) + .484ca\ cb\ sg^2\ cg^2 \\
&\quad - cg((.527ca + .484sa^2)cb + .484ca\ sb^2)sg^2 - (cb(.699ca - .527sa^2 + .484ca\ sa^2) + \\
&\quad .484ca\ cb\ sb^2 - (.527ca + .484sa^2)sb^2)sg^2 \\
U_{25} &= cg(-.484ca\ sb\ cb^3 + (.699sa + .527ca\ sa - .484ca^2\ sa)sb \\
&\quad + (.527ca + .484sa^2)cb^2\ sb + (.699ca - .527sa^2 + .484ca\ sa^2)cb\ sb) + .484ca\ cb\ cg^3\ sg^2 \\
&\quad - cg^2(cb(.527ca + .484sa^2) + .484ca\ sb^2)sg^2 - cg(cb(.699ca - .527sa^2 + .484ca\ sa^2) + \\
&\quad .484ca\ cb\ sb^2 - (.527ca + .484sa^2)sb^2)sg^2 - (cb(.699sa + .527ca\ sa - .484ca^2\ sa) + \\
&\quad .484ca\ cb^2\ sb^2 - cb(.527ca + .484sa^2)sb^2 + (.699ca - .527sa^2 + .484ca\ sa^2)sb^2)sg^2 \\
U_{26} &= -.484ca\ cb\ sg\ cg^4 + (-.484ca\ cb^3\ sb + (.699sa + .527ca\ sa - .484ca^2\ sa)sb \\
&\quad + cb^2(.527ca + .484sa^2)sb + cb(.699ca - .527sa^2 + .484ca\ sa^2)sb)sg \\
&\quad + cg^3((.527ca + .484sa^2)cb + .484ca\ sb^2)sg + \\
&\quad cg^2(cb(.699ca - .527sa^2 + .484ca\ sa^2) + .484ca\ cb\ sb^2 - (.527ca + .484sa^2)sb^2)sg \\
&\quad + cg(cb(.699sa + .527ca\ sa - .484ca^2\ sa) + .484ca\ cb^2\ sb^2 - cb(.527ca + .484sa^2)sb^2 \\
&\quad - (.699ca - .527sa^2 + .484ca\ sa^2)sb^2)sg. \tag{5}
\end{aligned}$$

3 $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ For equal sterile neutrino masses

Assuming that all three sterile neutrinos have the same mass, sterile-active neutrino mass differences are $\delta m_{4j}^2 = m_4^2 - m_j^2 \simeq .9(eV)^2$, with δm_{4j}^2 taken from the best fit to neutrino oscillation data[4] (see Ref[4] for references to earlier experiments), from Eq(3) $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ is

$$\begin{aligned} \mathcal{P}(\nu_\mu \rightarrow \nu_e) = & Re[U_{11}U_{21}[U_{11}U_{21} + U_{12}U_{22}e^{-i\delta L} + U_{13}U_{23}e^{-i\Delta L} + \\ & (U_{14}U_{24} + U_{15}U_{25} + U_{16}U_{26})e^{-i\gamma L}] + \\ & U_{12}U_{22}[U_{11}U_{21}e^{-i\delta L} + U_{12}U_{22} + U_{13}U_{23}e^{-i\Delta L} + \\ & (U_{14}U_{24} + U_{15}U_{25} + U_{16}U_{26})e^{-i\gamma L}] + \\ & U_{13}U_{23}[U_{11}U_{21}e^{-i\Delta L} + U_{12}U_{22}e^{-i\Delta L} \\ & + U_{13}U_{23} + (U_{14}U_{24} + U_{15}U_{25} + U_{16}U_{26})e^{-i\gamma L}] + \\ & (U_{14}U_{24} + U_{15}U_{25} + U_{16}U_{26})[(U_{11}U_{21} + U_{12}U_{22} \\ & + U_{13}U_{23})e^{-i\gamma L} + U_{14}U_{24} + U_{15}U_{25} + U_{16}U_{26}] , \end{aligned} \quad (6)$$

with $\delta = \delta m_{12}^2/2E$, $\Delta = \delta m_{13}^2/2E$, $\gamma = \delta m_{jk}^2/2E$ ($j=1,2,3; k=4,5,6$). The neutrino mass differences are $\delta m_{12}^2 = 7.6 \times 10^{-5}(eV)^2$, $\delta m_{13}^2 = 2.4 \times 10^{-3}(eV)^2$; and $\delta m_{jk}^2 (j = 1, 2, 3; k = 4, 5, 6) = 0.9(eV)^2$ [4].

From Eq(6)

$$\begin{aligned} \mathcal{P}(\nu_\mu \rightarrow \nu_e) = & U_{11}^2 U_{21}^2 + U_{12}^2 U_{22}^2 + U_{13}^2 U_{23}^2 + \\ & (U_{14}U_{24} + U_{15}U_{25} + U_{16}U_{26})^2 + \\ & 2U_{11}U_{21}U_{12}U_{22}\cos\delta L + \\ & 2(U_{11}U_{21}U_{13}U_{23} + U_{12}U_{22}U_{13}U_{23})\cos\Delta L + \\ & 2(U_{14}U_{24} + U_{15}U_{25} + U_{16}U_{26}) \\ & (U_{11}U_{21} + U_{12}U_{22} + U_{13}U_{23})\cos\gamma L . \end{aligned} \quad (7)$$

Note that $\alpha \simeq 9.2^\circ$ from a recent analysis of MiniBooNE data, which was used in a recent study of $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ with one sterile neutrino[8, 9]. The figure below shows $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ with $\alpha = \beta = \gamma = 0^\circ$, giving the results of a recent 3x3 S-matrix calculation[12]. In the figure, for the other curves, the sterile-active mixing angle $\alpha = 9.2^\circ$, while β and γ are chosen to be 9.2° and 20° to compare the 6x6 to the previous 3x3 results.

Using Eq(7), one finds $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ for the 6x6 vs 3x3 theories:




Figure 1: $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ for MINOS(L=735 km), MiniBooNE(L=500m), JHF-Kamioka(L=295 km), and CHOOZ(L=1.03 km). (a) solid, for $\alpha = \beta = \gamma = 9.2^\circ$; (b) dashed, for $\alpha, \beta, \gamma = 9.2^\circ, 20^\circ, 20^\circ$; (c) dash-dotted curve for $\alpha = \beta = \gamma = 0^\circ$ giving the 3x3 result .

4 Conclusions

From the figure we note that with the small mixing angle, $\alpha = \beta = \gamma = 9.2^\circ$, taken from the MiniBooNE analysis for s_α , for MINOS, MiniBooNE, and JHF-Kamioka there is significant difference between our 6x6 and the earlier 3x3 prediction for $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$, given by $\alpha = \beta = \gamma = 0^\circ$. For the larger 20° mixing angles for β and γ , which are not known, there is a much larger difference between the 6x6 and 3x3 theories for these three experimental set-ups. For CHOOZE, however, $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ is not significantly dependent on the mixing angles α, β, γ for the values used, and is similar to the 3x3 prediction.

There are many different choices for the parameters needed for this study, which we shall investigate in future work.

Acknowledgements

This work was carried out while LSK was a visitor at Los Alamos National Laboratory, Group P25. The author thanks Dr. William Louis for information about recent and future neutrino oscillation experiments, and Dr. Terrance Goldman for advice on the mixing angles.

References

- [1] E. Akhmedov, P. Huber, M. Linder, and T. Ohlsson, Nucl. Phys **B 608**, 394 (2001)
- [2] M. Jacobson and T. Ohlsson, Phys. Rev. **D 69**, 013003 (2004)
- [3] E.M. Henley, M.B. Johnson, and L.S. Kisslinger, arXiv:1102.5106/ hep-ph; Int. J. Mod. Phys. **E20**, 2453 (2011)
- [4] A.A. Aguilar-Arevalo *et. al.* (MiniBooNE Collaboration), Phys. Rev. Lett. **110**, 161801 (2013)
- [5] Joachim Kopp, Pedro A.N. Machado, Michele Maltoni, Thomas Schwetz, arXiv:1303.3011/hep-ph
- [6] S. Gariazzo, C. Giunti, M. Laveder, Y.F. Li, E.M. Zavanin, arXiv:1507.08204/hep-ph
- [7] Jiro Arafune and Joe Sato, Phys. Rev. **D 55**, 1653 (1997)
- [8] Leonard S. Kisslinger, arXiv:1309.4983/hep-ph; Int.J.Theor.Phys **53**, 3201 (2014)

- [9] Leonard S. Kisslinger, arXiv:1409.1610/hep-ph; Int.J.Theor.Phys **54**, 2141 (2015)
- [10] T. Goldman, Int. J. Mod. Phys. **A22**, 4967 (2007)
- [11] T. Goldman and G.J. Stephenson, Jr., Phys. Rev. **D 24**, 236 (1981)
- [12] Leonard S. Kisslinger, arXiv:1408.0310/hep-ph; Mod. Phys. Lett. A, **30**, 1550014 (2015)